



The Investigational Drugs database  
**DRUG report**

QuickSearch:

SEARCH

Drug

## troglitazone

Company

Sankyo Co Ltd

Highest Dev Status

Withdrawn

Indications

Non-insulin dependent diabetes  
Diabetes mellitus

Actions

Insulin sensitizer  
PPAR agonist  
Hypoglycemic agent  
5-HETE modulator

Browse Database

Browse Indexes

Search Center

IDdb Portal

Personal

IDeals

## Summary

**Troglitazone**, a peroxisome proliferator activated receptor (PPAR) agonist, was launched by Sankyo in Japan in April 1997, and in the US by Parke-Davis in March 1997 [270683] for the treatment of non-insulin dependent diabetes (NIDDM) [237320] but has since been withdrawn from these markets. Some preclinical data, reported in April 2000, described troglitazone as having potential as an anticancer agent in head and neck squamous cell carcinoma [362460].

In March 2000, Warner-Lambert voluntarily discontinued the sale of troglitazone from the US market. The company decided, following discussions with the FDA, that it was in the best interests of patients to discontinue marketing troglitazone at that time [360409]. The FDA subsequently opened an investigation into allegations that Warner-Lambert had manipulated adverse-event reports regarding the toxic liver effects of troglitazone [360696]. Sankyo also voluntarily discontinued the sale of troglitazone in both Japan and the US [363389]. By January 2003, the FDA had withdrawn its approval [476981].

In April 2000, class action lawsuits against Warner-Lambert and Parke-Davis were filed in the US District Court for the District of New Jersey and in the federal court in Philadelphia. The suits allege that the defendants misled potential users of troglitazone concerning the health risks associated with the drug and alleges that the defendant's product warnings were wholly inadequate and failed to warn prescribing physicians and patients of the actual heart and liver risks associated with the compound [362272].

Related information
COMPANY
REFERENCES
NEWS
PATENT
Actions
EMAIL IF UPDATED
ADD TO LIBRARY
PRINT VIEW
WORD VIEW
FIND SIMILAR

BEST AVAILABLE COPY

1/7 ~-~

SN 10/606, 632

file:///D:/DOCUMENT1/19940161/LOCALS1/Temp/GSBJ2S86.htm

2006/03/06



Results Summary > Drug Report

QuickSearch:

SEARCH  
Drug

- IDdb Alerts
- Browse Database
- Browse Indexes
- Search Center
- IDdb Portal
- Personal
- IDeas

pioglitazone

Company

Takeda Pharmaceutical Co Ltd

Highest Dev Status

Launched

Indications

Hypertlipidemia  
Hypertension  
Myocardial infarction  
Atherosclerosis  
Cerebrovascular Ischemia  
Non-insulin dependent diabetes

Actions

Insulin sensitizer  
PPAR gamma agonist  
Hypoglycemic agent  
Antypercholesterolemic agent  
Tablet formulation

Technologies

Reason for update on 23 February 2006

Minor editorial amendment

Summary

Pioglitazone is a peroxisome proliferator-activated receptor (PPAR)-gamma agonist, which, by late-2000, had been launched extensively by Takeda for the treatment of non-insulin dependent diabetes mellitus (NIDDM; type 2 diabetes) [181756]. Takeda was initially copromoting the drug in the US with Eli Lilly, and this was expanded to include more than 70 countries within Europe, the Middle East, Africa and Asia Pacific in August 1999 [338000]. The company is also developing the drug for the potential treatment of atherosclerosis, hypertension and for the potential prevention of myocardial infarction and stroke. In May 2003, it was in phase III trials for atherosclerosis [491975]; these trials were ongoing in March 2005 [606970], [515419]. In November 2004, clinical data showing the efficacy of the drug for the treatment of hypertension were published [570792], [570772].

ATHEROSCLEROSIS

In November 2004, clinical data demonstrating the relative efficacy of pioglitazone (30 mg/day) compared to rosiglitazone (4 mg/day) on blood lipids and insulin sensitivity were presented at the ADA meeting in New Orleans, LA. Results from the randomized, double-blind, multicenter trial in 735 patients with type 2 diabetes showed that both drugs exhibited similar degrees of glycemic control. However, at week 24,

Related information
COMPANY
REFERENCES
NEWS
PATENT
Actions
EMAIL IF UPDATED
ADD TO LIBRARY
PRINT VIEW
WORD VIEW
FIND SIMILAR



Results Summary > Drug Report

QuickSearch:

SEARCH  
Drug

- IDdb Alerts
- Browse Database
- Browse Indexes
- Search Center
- IDdb Portal
- Personal
- IDEALS

rosiglitazone

Company

SmithKline Beecham plc

Highest Dev Status

Launched

Indications

Alzheimer's disease  
Psoriasis  
Rheumatoid arthritis  
Asthma  
Ulcerative colitis  
Non-insulin dependent diabetes

Actions

Insulin sensitizer  
PPAR gamma agonist  
Anti-inflammatory  
Tablet formulation

Technologies

Reason for update on 09 February 2006

Minor editorial amendment, 1 reference added [649382]

## Summary

Rosiglitazone (Avandia), a peroxisome proliferator-activated receptor (PPAR)-gamma receptor agonist, has been developed and launched in several major markets by SmithKline Beecham (SB; now GlaxoSmithKline, GSK) as a treatment for non-insulin dependent diabetes mellitus (NIDDM/type II) [263572]. It is also in development for several other indications, including Alzheimer's disease (AD) and asthma [515870], [632116]. It was launched in the US and Mexico for diabetes in June 1999 [327686], [333230], and by June 2000, had also been launched in the UK and Germany [376124]. By December 2003, the compound was in phase II trials for Alzheimer's disease (AD) [515870] and in November 2005 trials were ongoing [638027]; by February 2006, the company expected an extended-release formulation to enter phase II trials in Alzheimer's disease later that year [649382]. In July 2005, a phase II trial was initiated in asthma patients, due to be completed by March 2006 [632116]. In November 2005, phase II trials had been initiated in rheumatoid arthritis (RA) [638027]. By February 2006, Glaxo expected phase III trials of rosiglitazone with simvastatin (qv) to begin later that year [649382].

The compound has also previously been developed for psoriasis: in 2003, phase III trials for psoriasis were initiated [515870]. However, the results reported in October 2004 showed that the compound had no efficacy for this indication [628420], [628419] and no further development for psoriasis has been reported

Related Information
COMPANY
REFERENCES
NEWS
PATENT
Actions
EMAIL IF UPDATED
ADD TO LIBRARY
PRINT VIEW
WORD VIEW
FIND SIMILAR

http://lddb3.com/lddb3/lddb3\_2/reports.display?id=8976&template=Drug&i\_query\_id=6571738

2006/03/06

Iddb3::Drug report

1/8 ページ



The Investigational Drugs database  
**DRUG report**

QuickSearch:

Drug Report

netoglitazone

Company

Highest Dev Status

Indications

Actions

Mitsubishi-Tokyo Pharmaceuticals Inc

Phase 2 Clinical

Metabolic disorder

Non-insulin dependent diabetes

Insulin sensitizer

PPAR gamma agonist

PPAR alpha agonist

Hypoglycemic agent

SEARCH  
Drug  
IDdb Alerts  
Browse Database  
Browse Indexes  
Search Center  
IDdb Portal  
Personal



Reason for update on 12 January 2006

Minor editorial amendment

## Summary

Perlegen and Mitsubishi Pharma (formerly Mitsubishi-Tokyo) are developing netoglitazone (MCC-555; RWJ-241947; Isaglitazone), a thiazolidinedione triple peroxisome proliferation activated receptor (PPAR) agonist (alpha, gamma and delta), as a potential treatment for type 2 diabetes and other metabolic diseases [204644], [423717], [594833]. By March 2004, phase II trials had been initiated in Japan [538366], [520195]; these trials were ongoing in November 2004 [595948]. In April 2005, Perlegen was planning US clinical trials [594833].

Johnson & Johnson (J&J) was previously developing the compound outside of Japan and, in April 1999, had listed netoglitazone as being in phase II trials [304271], [322655]; however, by May 2003, the agreement between Mitsubishi and J&J had been terminated [491163].

## PRECLINICAL DATA

In June 2003, preclinical data on netoglitazone were presented at the 63rd ADA meeting in New Orleans, LA. Netoglitazone inhibited TNFalpha-induced vascular cell adhesion molecule-1 (VCAM-1) expression in a dose-dependent manner in vitro. Monocyte chemotactic protein-1 (MCP-1) secretion from human aortic endothelial cells was dose-dependently inhibited by netoglitazone, suggesting beneficial effects on endothelium in early stage of atherosclerosis, mediated by PPAR alpha or PPAR delta [492285].

Related Information	
COMPANY	
REFERENCES	
NEWS	
PATENT	
Actions	
EMAIL IF UPDATED	
ADD TO LIBRARY	
PRINT VIEW	
WORD VIEW	
FIND SIMILAR	

http://iddb3.com/iddb3\_2/reports.display?id=17476&template=Drug&i\_query\_id=65716256i\_clear\_trail=Y

2006/03/06

iddb3::Drug report

1/9 ^-j



## Drug Report

## QuickSearch:

 SEARCH  
 Drug

reglitazar

Company

Japan Tobacco Inc

Highest Dev Status

No Development Reported

Indications

Non-insulin dependent diabetes

Actions

 Insulin sensitizer  
 PPAR gamma agonist  
 PPAR alpha agonist

IDdb Alerts

Browse Database

Browse Indexes

Search Center

IDdb Portal

Personal

Ideals

Reason for update on 24 March 2004

5 references added [515261, 389917, 431904, 473077, 464687], indexing updated, literature evaluation added, one or more development status entries have been updated

## Summary

Japan Tobacco was developing reglitazar (JTT-501), an insulin sensitizer and PPARalpha and gamma agonist, as a potential oral treatment for Type II diabetes [249397]. By 1998, reglitazar was in phase II trials in the UK for Type II diabetes [290700]. In July 2001, phase III trials were expected to commence later that year [402336]. However, in October 2002, the company decided to terminate development of the drug candidate based on an assessment of its results [466412].

Pharmacia Corp (formerly Pharmacia & Upjohn) had development and marketing rights worldwide except for Japan and Korea [289901], although in April 2002, the company reported that it was no longer involved in the development of reglitazar [445305]. Phase II trials had commenced in Japan, where Wellfide (now Mitsubishi Pharma) was codeveloping the compound [304419]; however, there has been no development reported on this collaboration since 1998.

In September 2001, preclinical data were presented at the European Association for the Study of Diabetes meeting, Glasgow, UK. In vitro and in vivo studies suggested that reglitazar exerted its antidiabetic action, following a pathway similar to that through which insulin exerts its own effect [429965].

In June 2001, data on reglitazar were presented at the 61st ADA meeting in Philadelphia, PA. These data demonstrated that, in the Type II db/db male mice model, EC50 values (micromolar) for PPARalpha and gamma agonism were (alpha/gamma): rosiglitazone (qv) 0.1/5; pioglitazone (qv) 1/7; NNC-61-0029 (qv) 0.6/3; reglitazar 0.4/2; MCC-555 (qv) 3/0.1; KRP-297 (qv) 0.5/0.4; GT-262570 (qv) 0.002/0.3; and, fenofibrate

Related information	
<a href="#">COMPANY</a>	
<a href="#">REFERENCES</a>	
<a href="#">NEWS</a>	
Actions	
<a href="#">EMAIL IF UPDATED</a>	
<a href="#">ADD TO LIBRARY</a>	
<a href="#">PRINT VIEW</a>	
<a href="#">WORD VIEW</a>	
<a href="#">FIND SIMILAR</a>	

[http://iddb3.com/iddb3/iddb3\\_2/reports.display?id=15328&template=Drug&i.query\\_id=6571621&i.clear\\_trail=Y](http://iddb3.com/iddb3/iddb3_2/reports.display?id=15328&template=Drug&i.query_id=6571621&i.clear_trail=Y)

2006/03/06

Iddb3::Drug report



QuickSearch:




[IDdb Alerts](#)  
[Browse Database](#)  
[Browse Indexes](#)  
[Search Center](#)  
[IDdb Portal](#)  
[Personal](#)

Drug Report

farglitazar

Company

Glaxo Wellcome plc

Highest Dev Status

Phase 2 Clinical

Indications

Unidentified

Fibrosis

Cardiovascular disease

Non-insulin dependent diabetes

Actions

Insulin sensitizer

PPAR gamma agonist

Retinoid X receptor modulator

Reason for update on 02 December 2005

one or more development status entries have been updated, indexing updated

#### Summary

GlaxoSmithKline (formerly Glaxo Wellcome) is developing farglitazar (GW-262570), a peroxisome proliferator-activated receptor (PPAR)-gamma agonist and retinoid X receptor modulator, for the potential treatment of hepatic fibrosis [638027], and investigating it for the potential treatment of cardiovascular diseases [470611], [464521]. In November 2005, farglitazar was in phase II trials for hepatic fibrosis [638027].

The compound was previously under development for type II diabetes, for which it reached phase III trials [730289], [335170], [399771], [409231]. However, in October 2001, it was reported that development of the compound for this indication had ceased as it did not meet its target profile. Alternative indications for the compound were being explored at that time, but the company would not disclose these, only revealing that the field would not be obvious to people who had not studied the group of compounds very closely [426569], [427414].

#### CARDIOVASCULAR DISEASE

In November 2002, preclinical data on farglitazar were presented at the AHA meeting in Chicago, IL. Rats receiving 8 mg/kg bid po farglitazar showed PPAR-gamma activation through detection of increased mRNA levels of the target genes FABP3 and aP2. Increased vasodilator NO levels ( $p < 0.05$ ) and fluid retention were observed, while the glomerular filtration rate, effective renal plasma flow and renal filtration fraction

Related information	
<a href="#">COMPANY</a>	
<a href="#">REFERENCES</a>	
<a href="#">NEWS</a>	
<a href="#">PATENT</a>	
<a href="#">Actions</a>	
<a href="#">EMAIL IF UPDATED</a>	
<a href="#">ADD TO LIBRARY</a>	
<a href="#">PRINT VIEW</a>	
<a href="#">WORD VIEW</a>	
<a href="#">FIND SIMILAR</a>	

[http://iddb3.com/iddb3/iddb3\\_2/reports.display?i\\_d=14652&template=Drug&i\\_query\\_id=6571659&i\\_clear\\_trail=Y](http://iddb3.com/iddb3/iddb3_2/reports.display?i_d=14652&template=Drug&i_query_id=6571659&i_clear_trail=Y)

2006/03/06

1/11 へ→



Drug Report

QuickSearch:

SEARCH  
Drug

- IDdb Alerts
- Browse Database
- Browse Indexes
- Search Center
- IDdb Portal
- Personal
- IDeals

YM-440

Company: **Astellas Pharma Europe BV**  
 Highest Dev Status: **No Development Reported**  
 Indications: **Non-insulin dependent diabetes**  
 Actions: **Insulin sensitizer**  
**Hypoglycemic agent**

Reason for update on 24 March 2004

1 reference added [390055], indexing updated, one or more development status entries have been updated, indexing updated, one or more development status entries have been updated

Summary

**Yamanouchi Europe** was developing **YM-440**, an insulin sensitizer, for the potential treatment for non insulin-dependent diabetes. The company had begun phase II trials in Europe by 1999 [326891]; however, the compound has not appeared on a company pipeline since May 2000 [365756], and no further development has been reported since. By May 2002, **YM-178** (qv), a beta 3 receptor agonist, appeared to have superseded **YM-440** in Europe for this indication.

An oral dose of **YM-440** 30 mg/kg od, for four days caused a reduction of 39% in blood glucose levels in male mice. Although similar in some respects to **troglitazone** (qv), Yamanouchi claims that **YM-440** has a different mechanism of action to the thiazolidinediones as a class. Specifically, **YM-440** showed only weak activity against PPAR-gamma [279360].

Administration of 100 mg/kg, **YM-440** to diabetic db/db mice for two weeks significantly decreased blood glucose concentrations (418 to 243 mg/dl); notably, there was no significant increase in body weight gain at the end of the treatment (in contrast to **troglitazone** and **pioglitazone**, (qv)). **YM-440** did not affect PPAR-gamma activity [293702].

**YM-440** is (Z)-1,4-bis-4-[(3,5-dioxo-1,2,4-oxadiazolidin-2-yl)-methyl]phenoxybut-2-ene [293702].

In February 1999, Lehman Brothers predicted the first major product launch to be in 2002, with sales

- Related Information
- COMPANY
- REFERENCES
- NEWS
- Actions
- EMAIL IF UPDATED
- ADD TO LIBRARY
- PRINT VIEW
- WORD VIEW
- FIND SIMILAR

iddb3::Drug report



QuickSearch:

SEARCH  
Drug

- IDdb Alerts
- Browse Database
- Browse Indexes
- Search Center
- IDdb Portal
- Personal



Drug Report

**KRP-297**  
Company  
Highest Dev Status  
Indications  
Actions  
Technologies  
Reason for update on 30 August 2005  
1 reference added [520284]

**Kyorin Pharmaceutical Co Ltd**  
Discontinued  
Hyperlipidemia  
Non-insulin dependent diabetes  
Insulin sensitizer  
PPAR gamma agonist  
PPAR alpha agonist  
Antihyperlipidemic agent  
Tablet formulation

Related Information	
COMPANY	
REFERENCES	
NEWS	
PATENT	
Actions	
EMAIL IF UPDATED	
ADD TO LIBRARY	
PRINT VIEW	
WORD VIEW	
FIND SIMILAR	

Summary

Merck & Co subsidiary Banyu, in collaboration with Kyorin, was developing KRP-297 (MK-767, L-410198), a PPAR alpha and gamma agonist, for the potential treatment of diabetes and hyperlipidemia [311944], [506609]. By September 2002, the companies were conducting phase II trials in Japan [488534], [516815]. Merck had initiated phase III trials in December 2002 [473816], but discontinued development in November 2003 for safety reasons [514245]. In December 2003, Kyorin announced that it was also discontinuing the development of KRP-297 [520055].

As of December 2002, Merck had been developing KRP-297 as MK-767 [473816]; however, in November 2003, the company discontinued its phase II development program because a long-term safety assessment program had identified a rare form of malignant tumors in mice [514245]. At this time, Kyorin was in discussion with Merck on the future development of KRP-297 in Japan [518196].

CLINICAL STUDIES

In March 2003, clinical data on KRP-297 were presented at the Lorenzini Foundation's Second International Meeting on PPARs in Florence, Italy. A double-blind, randomized, placebo-controlled, single-dose trial was performed to determine the safety, tolerability, pharmacokinetics and lipid effects of KRP-297. Single oral doses from 1 to 80 mg were well tolerated. Plasma AUC and Cmax values increased with dose and the terminal half-life of KRP-297 was 36 h. A standard breakfast did not affect the absorption of a 5 mg oral

http://iddb3.com/iddb3\_2/reports.display?id=21617&template=Drug&i\_query\_id=6571669&i\_clear\_trail=Y

2006/03/06

1/10 ページ





## Drug Report

## QuickSearch:

SEARCH

Drug

LY-282449

Company

Eli Lilly &amp; Co

Highest Dev Status

No Development Reported

Indications

Diabetic complication

Actions

Insulin sensitizer  
PPAR gamma agonist

IDdb Alerts

Related Information

COMPANY

REFERENCES

NEWS

Actions

EMAIL IF UPDATED

ADD TO LIBRARY

PRINT VIEW

WORD VIEW

FIND SIMILAR

Browse Database

Browse Indexes

Search Center

IDdb Portal

Personal



## Summary

TA-174 is a thiazolidinedione peroxisome proliferator-activated receptor (PPAR) gamma agonist which was under investigation by Eli Lilly in collaboration with Tanabe as a hypoglycemic agent for the potential treatment of diabetic complications. Lilly discontinued development of the drug in 1993 due to toxicity problems [155124] and although Tanabe continued to investigate the compound, no development has been reported by the company since 1997 [367139].

## Development Status

## HISTORY

## Detailed status for Eli Lilly &amp; Co

Indication	Country	Status	Reference	Date
Diabetic complication	US	Discontinued	155124	01 September 1993
Detailed status for Tanabe Seiyaku Co Ltd				
Indication	Country	Status	Reference	Date
Diabetic complication	Japan	No Development Reported		18 May 2000

[http://iddb3.com/iddb3/iddb3\\_2/reports.display?id=6053&template=Drug&i\\_query\\_id=6571675&i\\_clear\\_trail=Y](http://iddb3.com/iddb3/iddb3_2/reports.display?id=6053&template=Drug&i_query_id=6571675&i_clear_trail=Y)

2006/03/06

iddb3::Drug report



Drug Report

QuickSearch:

NC-2100

Company

Highest Dev Status

Indications

Actions

Nippon Chemiphar Co Ltd

Discovery

Non-insulin dependent diabetes

Insulin sensitizer

PPAR gamma agonist

Reason for update on 08 May 2003

1 reference added [488447]

[IDdb Alerts](#)  
[Browse Database](#)  
[Browse Indexes](#)  
[Search Center](#)  
[IDdb Portal](#)  
[Personal](#)  
[Ideals](#)

## Summary

NC-2100 is a thiazolidinedione PPAR-gamma activator which is under investigation by Nippon Chemiphar for the potential treatment of diabetes. In KKAY obese mice NC-2100, 0.1% for two weeks, lowered plasma glucose and triglyceride concentrations to levels comparable to those achieved with PPAR-gamma activators such as pioglitazone (qv) and troglitazone (qv) [371071].

## Development Status

Detailed status for Nippon Chemiphar Co Ltd

Indication	Country	Status	Reference	Date
Non-insulin dependent diabetes	Japan	Discovery	371071	16 June 2000

## Chemistry

Structure

[VIEW MOL FILE](#)

Related information
<a href="#">COMPANY</a>
<a href="#">REFERENCES</a>
<a href="#">NEWS</a>
Actions
<a href="#">EMAIL IF UPDATED</a>
<a href="#">ADD TO LIBRARY</a>
<a href="#">PRINT VIEW</a>
<a href="#">WORD VIEW</a>
<a href="#">FIND SIMILAR</a>

[http://iddb3.com/iddb3/iddb3\\_2/reports.display?id=29211&template=Drug&i\\_query\\_id=6571679&i\\_clear\\_trail=Y](http://iddb3.com/iddb3/iddb3_2/reports.display?id=29211&template=Drug&i_query_id=6571679&i_clear_trail=Y)

2006/03/06

1/6 1-2



Drug Report

QuickSearch:

SEARCH  
Drug

- IDdb Alerts
- Browse Database
- Browse Indexes
- Search Center
- IDdb Portal
- Personal
- IDeals

muraglitazar

Company

Bristol-Myers Squibb Co

Highest Dev Status

Pre-registration

Indications

Metabolic disorder  
Non-insulin dependent diabetes

Actions

Insulin sensitizer  
PPAR gamma agonist  
PPAR alpha agonist  
Hypoglycemic agent  
Antihyperlipidemic agent

Technologies

Oral formulation

Reason for update on 22 December 2005

one or more development status entries have been updated, 1 reference added [642539]

- Related information
- COMPANY
- REFERENCES
- NEWS
- PATENT
- Actions
- EMAIL IF UPDATED
- ADD TO LIBRARY
- PRINT VIEW
- WORD VIEW
- FIND SIMILAR

Summary

Bristol-Myers Squibb (BMS) is developing muraglitazar (BMS-298585; Pargluva; structure shown), a dual PPAR alpha/gamma agonist, as a potential oral treatment for type 2 diabetes and other metabolic disorders. An NDA was submitted in December 2004 [577532]. In October 2005, the FDA issued an approvable letter requesting additional information to support the cardiovascular safety profile of the drug [629049]. Later that month, BMS announced its would have to complete additional trials to secure approval. These were expected to take 5 years. At that time, BMS intended to meet with the FDA to discuss its options, including discontinuing development [630997]. By December 2005, discussions with the FDA were ongoing [642539].

Merck & Co was codeveloping the drug with BMS. However on receipt of the FDA approvable letter in October 2005, the companies began discussions to terminate the collaboration [630997]. In December 2005, BMS and Merck reached an agreement for the return of the drug to BMS [642539].

By March 2003, backup compounds (qv) were also under investigation [486672].

REGULATORY INFORMATION

[http://iddb3.com/iddb3/iddb3\\_2/reports.display?id=33451&template=Drug&i\\_query\\_id=6571694&i\\_clear\\_trail=Y](http://iddb3.com/iddb3/iddb3_2/reports.display?id=33451&template=Drug&i_query_id=6571694&i_clear_trail=Y)

2006/03/06

Iddb3: Drug report

1/15 〰—シ



QuickSearch:

Drug Report

tesaglitazar

Company

Highest Dev Status

Indications


Actions

Technologies

Reason for update on 10 February 2006

1 reference added [648528]. Minor editorial amendment

Summary



AstraZeneca plc

Phase 3 Clinical

Heart arrhythmia

Lipid metabolism disorder

Insulin dependent diabetes

Non-insulin dependent diabetes

PPAR gamma agonist

PPAR alpha agonist

Hypoglycemic agent

Antihypertensive agent

Oral formulation

Related Information

COMPANY

REFERENCES

NEWS

PATENT

Actions

EMAIL IF UPDATED

ADD TO LIBRARY

PRINT VIEW

WORD VIEW

FIND SIMILAR

**AstraZeneca** (formerly **Astra**) is developing **tesaglitazar** (AZ-242, AR-H039242, Gallda; structure shown), an oral dual PPAR alpha/gamma agonist, for the potential improvement of dyslipidemia and glyceric control in type 2 diabetes [275466], [377656]. In October 2003, phase III trials were being initiated [507400], [507401]; these were ongoing in October 2004. At that time, the anticipated filing date was moved from 2006 to 2007 as a result of AstraZeneca agreeing to extend long-term follow-up studies to 2 years following a worldwide regulatory authority review of the safety and toxicology of PPAR agonists [563036]. In November 2005, phase II trials were underway in Japan [638194]. By September 2005, tesaglitazar was on track for a 2007 submission [624516]; in February 2006, the expected submission date of the second half of 2007 was dependent upon the results of ongoing phase III studies and discussions with the FDA [648480], [648528].

In December 1999, the compound was also being developed as a potential antiarrhythmic drug [349551], [314472], [377656]; as of March 2001, however, diabetes and insulin resistance were the only indications listed on the company's pipeline [402040].

[http://iddb3.com/iddb3\\_2/reports.display?id=17061&template=Drug&i\\_query\\_id=6571699&i\\_clear\\_trail=Y](http://iddb3.com/iddb3_2/reports.display?id=17061&template=Drug&i_query_id=6571699&i_clear_trail=Y)

2006/03/06



Drug Report

QuickSearch:

SEARCH  
Drug

- IDdb Alerts
- Browse Database
- Browse Indexes
- Search Center
- IDdb Portal
- Personal
- IDeals

ragaglitazar

Company

Highest Dev Status

Indications

Actions

Dr Reddys Research Foundation

Discontinued

Hyperlipidemia

Non-insulin dependent diabetes

Insulin sensitizer

PPAR gamma agonist

PPAR alpha agonist

Hypoglycemic agent

Antihyperlipidemic agent

Reason for update on 05 May 2004

Minor editorial amendment, one or more development status entries have been updated, literature evaluation added, indexing updated

Summary

Ragaglitazar (NN-622, (-)-DRF-2725; structure shown), a PPAR (peroxisome proliferator-activated receptor) alpha and gamma agonist that regulates blood glucose and lipid levels, was being developed by Novo Nordisk (previously in collaboration with Novartis), under license from Dr Reddy, for the potential treatment of non-insulin dependent (Type II) diabetes [325344], [416231]. In November 2001, phase III trials were initiated both in the US and in Europe [428278], [440091]. In February 2002, these phase III trials were ongoing but Novo Nordisk revealed that it was planning to outlicense the compound [443186], [438928]. In July 2002, Novo Nordisk decided to suspend all current clinical trials of ragaglitazar and postponed all planned future trials. This decision was made in response to preclinical studies that revealed the formation of bladder tumors in one mouse and a number of rats after treatment with the drug. At this time, the company planned to continue all other activities in the development program until it completed a renewed benefit/risk assessment of ragaglitazar, the results of which would be ready by the first quarter of 2003. Filing for approval was expected to be delayed by two years [458773], [458774], [460379]. However, by February 2003, Novo Nordisk had decided not to pursue further development of the program [478217], [478168].

CLINICAL INFORMATION

Worldwide phase II trials had begun by September 2000 [381826], [385666]. By July 2001, Novo Nordisk expected to complete phase II trials later in that year [416231]; and, by September 2001, the

Related Information	
COMPANY	
REFERENCES	
NEWS	
PATENT	
Actions	
EMAIL IF UPDATED	
ADD TO LIBRARY	
PRINT VIEW	
WORD VIEW	
FIND SIMILAR	

http://iddb3.com/iddb3\_2/reports.display?id=23499&template=Drug&i\_query\_id=6571705&i\_clear\_trail=Y

2006/03/06

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: \_\_\_\_\_**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**